



Natureza & Conservação

Brazilian Journal of Nature Conservation

Supported by Boticário Group Foundation for Nature Protection

<http://www.naturezaeconservacao.com.br>



Essays and Perspectives

Grassland degradation and restoration: a conceptual framework of stages and thresholds illustrated by southern Brazilian grasslands



Bianca O. Andrade^{a,*¹}, Christiane Koch^{a,b,1}, Ilisi I. Boldrini^{a,c}, Eduardo Vélez-Martin^d, Heinrich Hasenack^e, Julia-Maria Hermann^b, Johannes Kollmann^b, Valério D. Pillar^{d,e}, Gerhard E. Overbeck^{a,c,d}

^a Graduate Program in Botany, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

^b Restoration Ecology, School of Life Sciences Weihenstephan, Technische Universität München, Freising, Germany

^c Department of Botany, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

^d Graduate Program in Ecology, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

^e Department of Ecology, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

ARTICLE INFO

Article history:

Received 24 August 2015

Accepted 27 August 2015

Available online 21 September 2015

ABSTRACT

Land degradation is a complex concept that integrates different aspects, including changes in soil conditions, biodiversity, productivity and socio-economic implications, compared to a reference state. We propose a new conceptual framework to analyze degradation stages and restoration thresholds in species-rich natural grasslands. The framework integrates different degradation stages with their respective thresholds and describes key processes of land-use change that lead to certain stages and thresholds. Specifically, we discuss two scenarios of grassland degradation, i.e. unsuitable grassland management and complete change of land use, sometimes followed by spontaneous recovery. We illustrate the framework with the case of south Brazilian grasslands, which are rich in biodiversity, but suffer from a series of degradation processes and are poorly considered from a conservation perspective. The conceptual framework can be applied by studies on degradation and restorability of tropical and subtropical grasslands after changes in management or transition to other land use; it will facilitate decisions on alternative management and conservation.

© 2015 Associação Brasileira de Ciência Ecológica e Conservação. Published by Elsevier Editora Ltda. All rights reserved.

Keywords:

Conservation

Grazing

Land-use change

Restoration

Biodiversity

Abiotic characteristics

* Corresponding author.

E-mail address: andradebo@gmail.com (B.O. Andrade).

¹ These authors contributed equally to this article.

<http://dx.doi.org/10.1016/j.ncon.2015.08.002>

1679-0073/© 2015 Associação Brasileira de Ciência Ecológica e Conservação. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Land-use change and degradation of natural ecosystems are principal causes for losses of biodiversity and ecosystem functions (Sala et al., 2000). Concrete numbers on land degradation are often only available for the complete loss of natural ecosystems, for example after conversion to more productive systems such as arable fields or tree plantations (for Brazil e.g. IBGE, 2012). However, degradation also includes less marked changes in structure, composition or ecological processes within the ecosystem. Therefore, degradation needs to be analyzed at different spatial scales, from a local focus on specific degradation processes to landscape and regional scales, and using diverse methods including remote sensing, plot-based measurements, experiments, expert knowledge and assessment of stakeholder experience (Reed et al., 2011). To provide a better understanding of the dynamics of degraded ecosystems and to facilitate mitigation of degradation processes and restoration, the concept of thresholds between alternative stages has been developed. While the framework suggested by Briske et al. (2006) focuses on definition and description of thresholds, Hobbs et al. (2009) described 'historical', 'hybrid' and 'novel' stages and identified 'restoration thresholds' when ecological, cultural or technical obstacles prevent a system to return to its original state. However, a synthesis of these approaches is missing so far.

Grasslands are among the ecosystems with highest species richness in the world (Wilson et al., 2012), and they provide a wide range of ecosystem services. Grasslands play an important role within the global carbon cycle, as 90% of their biomass is belowground, accumulation rates are high, and decomposition of organic material slow (Gibson, 2009). As main forage resource for livestock, grasslands are important for human well-being in many regions. They facilitate infiltration of water into the soil and thus to the maintenance of hydrological cycles. Finally, grasslands contribute to the landscape beauty of many regions. Thus, they are multi-functional systems but at the same time subjected to unsustainable use and conflicting interests.

Conceptual frameworks of grassland degradation have been developed specifically for rangelands, i.e. grazed systems, where stocking rates often are inadequate (e.g. Bestelmeyer, 2006). At the same time, large areas of grasslands are endangered due to land-use changes or have already been lost (Sala et al., 2000), especially in tropical and subtropical regions (Bond and Parr, 2010). Elsewhere, cropland on sites that originally had been grassland is abandoned for economic reasons, and grassland re-assembles spontaneously under re-introduction of moderate grazing management (Hölzel et al., 2002; Öster et al., 2009). This latter scenario, to our knowledge, has not yet been integrated in grassland or rangeland degradation models.

In this paper, we present a conceptual framework of grassland degradation that for the first time integrates different degradation stages and two types of recovery thresholds, including the most important processes related to land-use history causing stage transitions. We illustrate the framework with studies from grasslands in Rio Grande do Sul

(hereafter: RS) State, southern Brazil, where biodiversity is well studied, but where conservation of grassland has been neglected, and degradation processes are poorly studied (Overbeck et al., 2007, 2013). Finally, we point out how current limitations regarding knowledge on degradation processes and conservation state of grasslands should guide future research.

Conceptual framework of grassland degradation and restoration

The starting point of our framework is the use of traditionally managed grasslands as reference systems. This reference stage is comprised by grasslands composed of native species, with a specific and high biodiversity and considerable conservation value, usually due to extensive grazing and associated management practices (e.g. fire, mowing). The importance of management for maintenance of these systems and their diversity has been shown before (e.g. Overbeck et al., 2007; Pillar and Vélez, 2010 for southern Brazil).

As in Hobbs et al. (2009), changes of ecosystem properties can be displayed as biotic and abiotic changes at the local scale along two axes (Fig. 1). As biotic changes we consider deviation in species composition, vegetation structure (vertical and horizontal structural variables), basic ecological processes (e.g. pollination, seed dispersal), resulting from altered management, species introduction or conversion of land use. This includes alien species invasion. As abiotic changes we understand alteration in soil chemical and physical properties, as caused by fertilization or soil cultivation. The applicability of the framework is directly related to the quantification of these factors in case studies and the establishment of the thresholds according to changes in biotic and abiotic conditions in grasslands. It allows us to identify the stage of degradation of a given area and its potential for restoration. Changes in grassland management will cause properties of the system to change gradually, resulting in a decrease in resilience, but with resumption of the historical management the original properties might be reached again in many cases, i.e. a self-recovery threshold is not crossed (Fig. 1a, Scenario 1). If the grassland is converted to other land use, this will lead to an almost entire change of the original properties, e.g. complete loss of the aboveground plant community (Fig. 1b, Scenario 2), while abandonment of these land uses initiates re-colonization from the regional species pool (Fig. 1c). This process is of particular importance in regions of the world where grassland restoration techniques have not been developed. In this context, it needs to be recognized that spontaneous recovery may often mean colonization of non-typical or alien species, i.e. trajectories of community development that not necessarily lead to the desired state.

In the framework, the distinction between two types of thresholds is important: (1) a self-recovery threshold that describes until which point a recovery without additional technical measures is possible, and (2) a restoration threshold that identifies until which point an area can be restored with technical assistance (e.g. modification of soil




Fig. 1 – Conceptual framework on degradation and restoration of species-rich natural grasslands: (a) properties of the reference grassland are moderately altered if grassland management is changed, but modification is reversible; (b) after conversion to other land uses, the properties of the reference grassland are radically changed and ecosystem resilience is lost; (c) after abandonment some properties might recover ('self-recovery threshold') or be restorable ('restoration threshold'); and (d) integration of the various scenarios.

features, control of undesired species, introduction of desired species).

South Brazilian native grasslands: origin and development

In RS, grasslands occur in the highlands in the north of the state, where they form mosaics with Araucaria forest, and in its southern half, in the Pampa biome, where they dominate the landscape (Overbeck et al., 2007), continuing in Uruguay and Argentina (Fig. 2). South Brazilian grasslands are particularly rich in plant species, with about 2200 grassland plant taxa known only for RS (Boldrini, 2009).

Grasslands in the region are primary. They are relicts from cooler and drier periods, and were affected by forest expansion since approximately 5000 years BP, favoured by warmer and more humid climate, with increasing rates since 1500 years BP (Behling, 2002). After the extinction of large herbivores (Lima-Ribeiro and Diniz-Filho, 2013), these grasslands had been

maintained by anthropogenic fires and by grazing of small mammals (Cione et al., 2003; Behling and Pillar, 2007), and since the 17th century by introduced livestock. Today, beef production is an important economic activity in the region, with native plant species constituting most grassland vegetation. Available data indicate that plant diversity and forage production are highest under intermediate levels of grazing and intermediate fire frequency (Overbeck et al., 2005; Nabinger et al., 2009). Management thus can be considered essential for preservation of grassland biodiversity, as observed in many 'old-growth grasslands' (Veldman et al., 2015). However, transformation rates are high: between 1986 and 2002, grassland areas suffered losses of 16%, which corresponds to a loss rate of 1000 km² per year (Cordeiro and Hasenack, 2009).

Land-use change in the grassland region according to remote sensing

Evaluation of LANDSAT data shows that ca. 60% (104,553 km²) of former grassland area in southern Brazil had been




Fig. 2 – Location of the region used as an example for the proposed degradation framework, grasslands in Rio Grande do Sul, southern Brazil. Shown is the original distribution of natural grasslands in southeastern South America.

destroyed by 2002 (Fig. 3), mostly due to the conversion to arable fields or alien tree plantations. Losses of native grassland have not been uniform in space, but reflect soil properties and topographic constraints. In the Central Western Plateau region, native grasslands were nearly completely transformed into cropland, mostly for soybean. The coastal region has seen high rates of transformation, principally to rice and pine plantations. In the Northeastern Plateau, where soils are shallower, tree plantations and arable fields are the main causes of grassland losses. Here, land-use change has increased considerably within the past decade.

In the different regions of the state, 5–17% of the grassland area is classified as ‘degraded’, i.e. remote sensing data indicate former agricultural activity (e.g. tillage lines). Returning these areas to high nature conservation status might compensate for some of the ongoing losses through grassland conversion. However, compositional characteristics of vegetation itself (e.g. presence of alien species) cannot be observed by remote sensing data at this scale. In the Central Depression and in the Southwestern Grasslands of RS, for instance, a considerable proportion of the remaining grasslands mapped as ‘conserved’ actually have been degraded by alien forage species, which were deliberately seeded in some areas and colonized others, or by other processes. On-site ground studies are indispensable to verify the level of degradation and existence of self-recovery and restoration thresholds; remote

sensing data alone can only give a limited picture of grassland degradation.

Scenario 1: Degradation of grasslands after changes in management, and potential for self-recovery

Grazing

Most studies evaluating effects of different grazing intensities in RS focus on effects on forage or beef production (e.g. Maraschin and Corrêa, 1994; Moojen and Maraschin, 2002; Pinto et al., 2008), and only few analyze effects on species composition (Boldrini and Eggers, 1997; Soares et al., 2011) or soil properties (Bertol et al., 1998). Usually, grazed grasslands with moderate grazing intensity are formed by mosaics of intensively grazed patches dominated by prostrate grasses (e.g. *Axonopus affinis* Chase, *Paspalum notatum* Flüggé), and less grazed patches dominated by tussock grasses, small shrubs or other species less attractive for grazing animals (Boldrini and Eggers, 1997; Diaz et al., 2007). This heterogeneity of the vegetation leads to structural complexity and diversity.

If grazing is excluded – until now, a common practice in conservation units of Rio Grande do Sul – grassland structure quickly changes: tall tussock grasses, e.g. *Andropogon*




Fig. 3 – Distribution of grassland remnants and degraded grassland in RS State. The map is based on Landsat ETM+ images (spatial resolution: 30 m). Grassland areas with clearly visible signs of former land-use change (e.g. use as agricultural field) are considered as degraded, i.e. reflect past land-use change. Original grassland areas that have been completely transformed to other uses and not recovered to grassland comprise the category ‘converted’. Regions in white are those where natural vegetation cover is forest.

and *Sorghastrum* spp. (Boldrini and Eggers, 1997), become dominant, litter accumulates and microclimate at soil surface changes (Pallarés et al., 2005), drastically reducing plant species richness (Overbeck et al., 2005). In northeastern RS, the encroachment of shrubs of the genus *Baccharis* (principally, *B. uncinella* DC) and a slow invasion of forest pioneer species have been observed after abandonment (Oliveira and Pilar, 2004) despite the accumulation of grass biomass that may hinder fast recruitment of woody pioneers.

Overgrazing, on the other hand, can lead to the replacement of productive forage plants by species with lower forage quality, resulting in increasing cover of ruderal species and bare soil, while the contribution of highly nutritional C₃ grasses decreases (Pallarés et al., 2005). Ecosystem functions like water infiltration can be negatively affected as soil bulk density increases (Bertol et al., 1998). Either situation, when grazing is excluded or when the grassland is overgrazed, may be considered degraded due to changes in biotic and abiotic characteristics and the reduction of ecosystem resilience.

Introduction of alien C₃ and C₄ species and fertilization

Overseeding of natural grasslands with introduced alien species, often combined with fertilization and liming, aims to increase forage quality and quantity especially in winter (Nabinger et al., 2000). Although many native species have high productivity and nutritive potential, they are not available on the seed market, and introduced species are used instead; common species are *Lolium multiflorum* Lam., along with some European Fabaceae, e.g. *Trifolium repens* L. It has been shown that forage yield increases linearly with nitrogen addition (Brambilla et al., 2012) and promotes the increase of animal live weight gain per area, but also leads to marked changes in the floristic composition (Pallarés et al., 2005; Brambilla et al., 2012).

Introduced alien grassland species may become a serious problem when they spread to natural ecosystems. Large-scale invasion of these European-origin species that are used as forage, however, has not been reported yet, in contrast to some other species from temperate climates (e.g. *Ulex europaeus* L.) and, principally, some C₄ grass species of African

origin. For example, *Eragrostis plana* Nees had invaded up to 20% of Rio Grande do Sul's grassland area by the year 2008, causing severe reduction in forage quality and native plant diversity (Medeiros et al., 2004). Medeiros and Ferreira (2011) succeeded in at least partial suppression of populations of the invasive species by a combination of soil cultivation and seeding of both native and non-native forage species.

Fire

Fire has shaped the southern Brazilian grasslands during the past millennia (Behling and Pillar, 2007). The use of burns as management tools for livestock production, traditionally applied in the highland grasslands, is controversial due to concerns regarding possible negative impacts, and thus fire had been prohibited by state legislation until recently. Regular fires select for different species groups compared to grazing (tussock vs. prostrated grasses, respectively), and selectively affect some species groups (e.g. C₃ grasses when burns occur in winter), but do not seem to cause reductions in grassland diversity (Overbeck et al., 2005). Fidelis et al. (2012) showed higher species richness in frequently burned grassland plots in comparison to sites where burning and grazing had been excluded for some years. In fact, a large part of the grassland species is adapted to fire and can resprout from underground storage organs (Fidelis et al., 2014). Invertebrate communities and processes determined by their activity, e.g. decomposition, return to pre-fire values after relatively short periods, at least under patchy fire at fine scale (Podgaiski et al., 2013, 2014). Exclusion of fire in ungrazed areas, a common practice in conservation units, leads to the accumulation of dead biomass and the risk of high-intensity fires increases.

Potential for self-recovery and restoration after management-induced degradation

Potential for self-recovery should be high if the species present are members of the characteristic species pool of the region and changes in composition and structure were provoked mainly by changes in management. When soil conditions were changed, e.g. by liming or fertilization, self-recovery may be more difficult or even impossible, because introduced, sometimes invasive, species with high competitive abilities may be able to maintain high cover. Current research suggests that prescribed fire could be a conservation tool in areas where grazing is not feasible (Overbeck et al., 2005; Fidelis and Blanco, 2014), e.g. in conservation units. Thus, intensity and frequency of management are key factors for grassland recovery after any kind of degradation (Winter et al., 2012).

Scenario 2: Degradation and potential for self-recovery after complete conversion of grassland

For southern Brazil, some studies on ecosystem properties and ecological processes under different types of land use in former grassland areas are available. Table 1 synthesizes the

available data, considering variables of importance for regeneration or restoration after the end of intensive land use (e.g. seed bank) or that may persist in a changed condition over long periods of time.

Arable land use and its effects

Studies on the seed bank of arable fields on former grasslands in southern Brazil show that the number of grassland species decreases with management intensity, giving place to native or alien ruderal species (Favreto et al., 2007), thus reducing recovery potential of grassland. These results are in line with studies from other grassland ecosystems around the world that show higher abundance of weed species in the seed bank after agricultural use (Hutchings and Booth, 1996; Kiehl and Pfadenhauer, 2007). As vegetative recovery is the principal regeneration strategy of South Brazilian grasslands after disturbance (Fidelis et al., 2009), seed input from external sources as well as abiotic conditions should be limiting for recovery of the former grassland community: the bud bank likely does not persist through periods of intensive agricultural use.

Grassland conversion also results in changes of soil properties. Arable land use increases nutrient levels of the soil (Rheinheimer et al., 1998; Perin et al., 2003), leading to different trajectories of vegetation recovery. It is well known that large quantities of carbon stored in grasslands may be rapidly transferred to the atmosphere and lost when the grassland is plowed and converted to arable fields (Sala and Paruelo, 1997; Pillar et al., 2012). In a worldwide meta-analysis of carbon changes due to land-use changes, Guo and Gifford (2002) showed that a conversion of grasslands to crop rotation leads to a loss of 60% of belowground carbon. For southern Brazil, a decrease in C-stock in soils under conventional-tilling has been shown, with magnitude depending on management intensity; no-tillage systems result in much lower losses of C in soils (Bertol et al., 2004; Diekow et al., 2005).

Tree plantations

By 2009, 6000 km² (10%) of grasslands in RS were converted to plantations of pines, eucalypts or acacias (Gautreau and Velez, 2011). Native grassland vegetation composition is drastically affected under tree plantations, even with reduced soil disturbance for tree planting (Pillar et al., 2002). Gonçalves et al. (2008) found relatively low species richness and dominance of a few ruderals and some alien species like the grass *Melinis minutiflora* P. Beauv. in the soil seed bank under tree plantations in the Central Brazilian cerrado. Likewise, and in analogy to former agricultural fields, we can expect a low contribution of the seed bank in vegetation recovery.

Studies on effects of tree plantations on grassland soils give variable results (Table 1). Guo and Gifford (2002) stated that the conversion to plantations leads to a significant reduction of soil C stocks when coniferous species were used, while the effect with broadleaf species like eucalypts was not significant. For RS, Wiesmeier et al. (2009) found lower C stocks under pine plantations, while Mafra et al. (2008) could not show any changes. A growing number of literature examining potential for carbon sequestration in plantations is available, with changes based on the shift from belowground biomass

Table 1 – Review of studies on the effects of land-use change on ecosystem processes in grasslands of southern Brazil (RS, Santa Catarina, Paraná). All trends in comparison to reference grasslands (-, no studies available).

	Conversion to arable land	Conversion to forest plantation
Aboveground vegetation	No-tillage systems: more alien plant species ¹ After abandonment: lower floristic diversity, dominance of ruderal species ² or alien species	Usually no understory ³
Seed bank	Lower abundance and diversity of native plant species; higher abundance of ruderals and alien plant species ^{4,5}	-
Litter thickness and quality	-	-
C-stock and cycling	C stock better preserved under no-tillage ^{7,13} C stock reduced under conventional tilling by 22% ¹⁵	C stocks lower than in pasture, ⁷ or unchanged ⁹ Rio de la Plata grasslands (further to the South): tree plantation under high precipitation (level of RS) have reduced carbon stocks in soil when compared to grassland ¹² Pine: pH lower ^{7,8} or unchanged ⁹ ; N _{tot} falling ⁷ ; P lower or higher; K ⁺ lower; Al ³⁺ higher or unchanged ⁹ Eucalyptus: pH lower, K ⁺ , Ca ²⁺ , Mg ²⁺ lower, increase in Na ⁺ and Al ³⁺¹⁰
Soil pH and nutrient status	Increased nutrient load in topsoil ¹¹ ; pH raised ¹⁴ or lowered ¹¹	Pine spp.: soil density unchanged
Soil physical properties	Aggregate stability better preserved under no-tillage ⁶	

¹Favreto et al. (2007); ²Boldrini and Eggers (1997); ³Souza et al. (2013); ⁴Favreto and Medeiros (2006); ⁵Maia et al. (2008); ⁶Bertol et al. (2004); ⁷Wiesmeier et al. (2009); ⁸Schumacher et al. (2008); ⁹Mafra et al. (2008); ¹⁰Céspedes-Payret et al. (2012); ¹¹Rheinheimer et al. (1998); ¹²Berthrong et al. (2012); ¹³Pillar et al. (2012); ¹⁴Almeida et al. (2005); and ¹⁵Diekow et al. (2005).

dominance (grassland) to aboveground biomass with litter accumulation (plantation) (Guo et al., 2008). It has been shown that this potential strongly depends on soil types (Zinn et al., 2002), and might not be true for regions with high precipitation like RS, for which a decrease in soil carbon was observed (Guo and Gifford, 2002; Berthrong et al., 2012).

Potential for self-recovery and recovery thresholds after land-use change

Observational data indicate that the type of vegetation that develops after logging and abandonment differs considerably from that of reference grassland, and that species introduction likely is important if the objective is to restore grasslands. Zaloomis and Bond (2011) showed that species composition of grasslands established after logging of *Pinus elliottii* plantations in South Africa was markedly changed in comparison to reference grasslands. To what extent spontaneous regeneration will allow for return to pre-disturbance conditions is currently unknown and likely will depend on the time period of other land use (and with this, to what extent seeds or underground organs of target species remain in the ground), intensity of modifications of the site conditions (e.g. fertilization), and the landscape context (i.e. propagule sources). Both biotic and abiotic thresholds for recovery may exist, making active restoration necessary. As grasslands depend on management, intensity and/or frequency of grazing or fire will probably be crucial for the restoration process.

Discussion and conclusions

Land degradation studied on a regional scale often only considers conversion or complete losses of natural ecosystems, with limits to detect e.g. compositional changes, while at the local scale, finer effects of land management have to be

considered. Both perspectives are necessary for an improved assessment of degradation and potential restoration, and differences between the two types of degradation likely imply different perceptions regarding degradation, conservation or restoration.

For our study system, conclusive evidence is available that management is necessary for maintenance of diverse and productive grasslands in this region (Overbeck et al., 2007), and biodiversity conservation and livestock production can be considered as complementary management goals, allowing for sustainable use (e.g. Nabinger et al., 2009). Fire and grazing are selective forces that cause changes in grassland composition and structure, but their effects depend on frequency and intensity – both can contribute to conservation of biodiversity and productivity, but they can also be detrimental when frequency or intensity are too high or too low. A systematic and large-scale quantification of effects of different management types (especially intensification, Scenario 1) on different properties of the grasslands in the region is still missing, making it difficult to define degradation more precisely. Furthermore, the necessity of an integrated perspective of biodiversity conservation and sustainable use still is not widely recognized in the debate on conservation strategies (Overbeck et al., 2007; Pillar and Vélez, 2010).

Even though a considerable proportion of natural grasslands in southern Brazil have been converted to other land use (Scenario 2), concerns on potential restoration of these areas have been raised only recently (Overbeck et al., 2013), and are affecting the agenda of conservation authorities. Once supported by additional empirical data, our conceptual framework can support decision making and priority setting in nature conservation by identifying whether costly restoration measures will be necessary or adaptation in management could be sufficient for self-recovery. In this, it is important to recognize that not only biotic and abiotic characteristics

are covered by the framework, but that these can also be interpreted in terms of ecosystem functions and services (e.g. carbon sequestration, forage production).

Bestelmeyer (2006) points out problems and risks associated with threshold models: for instance, no single predictive thresholds – which would greatly facilitate management decisions – should be expected to exist, and parameters may reflect measurability, and not long-term degradation processes. Threshold models may become ‘insidious’ (**Bestelmeyer, 2006**), if they lead to the belief that certain areas are not restorable anymore, because some original features of the system cannot be recovered. This, however, is not a consequence of the model per se, but of a failure of recognizing the full range of features, processes and services of any type of ecosystem. The current debate on novel ecosystems (**Hobbs et al., 2013**) is centering exactly on the question of how to deal with systems that cannot be brought back to their original state. A conceptual framework of degradation and restoration based on a variety of biotic and abiotic variables (and their interactions, e.g. soil–plant feedbacks, e.g. **Suding et al., 2013**) has the potential to include different functions and services, and can contribute to a broader understanding of landscapes as multifunctional systems.

Young (2000) suggested that ecological restoration is the future of biological conservation. This means that degraded systems need to become a research focus, even in megadiverse countries where knowledge on biodiversity and functioning of natural ecosystems still has priorities. Conceptual frameworks such as the one presented here can support the study of degradation, conservation and restoration, in southern Brazil and elsewhere. In order to assess conservation or degradation state and restoration potential of degraded systems and to define the recovery and restoration thresholds, it is necessary to collect data on the full gradient from conserved to degraded systems, on a regional scale. For this, we need to develop rapid assessment methods of different parameters of the system, including abiotic and biotic variables as well as measures for ecosystem functions, as proposed by **Meyer et al. (2015)**.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

This contribution results from a CNPq-DFG cooperation project (490069/2011-8 and KO1741/2-1, respectively). CK receives support by the Evangelisches Studienwerk Villigst. BA by CAPES. EV, IB and VP receive support from CNPq.

REFERENCES

- Almeida, J.A., et al., 2005. Propriedades químicas de um Cambissolo Húmico sob preparo convencional e semeadura direta após seis anos de cultivo. *Rev. Bras. Ciênc. Solo* 29, 437–445.
- Behling, H., 2002. South and southeast Brazilian grasslands during Late Quaternary times: a synthesis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 177, 19–27.
- Behling, H., Pillar, V.D., 2007. Late quaternary vegetation, biodiversity and fire dynamics on the southern Brazilian highland and their implication for conservation and management of modern Araucaria forest and grassland ecosystems. *Philos. Trans. R. Soc. B* 362, 243–251.
- Berthrong, S.T., et al., 2012. Soil C and N changes with afforestation of grasslands across gradients of precipitation and plantation age. *Ecol. Appl.* 22, 76–86.
- Bertol, I., et al., 1998. Propriedades físicas do solo relacionadas a diferentes níveis de oferta de forragem numa pastagem natural. *Pesq. Agropec. Brasil.* 33, 779–786.
- Bertol, I., et al., 2004. Propriedades físicas do solo sob preparamento convencional e semeadura direta em rotação e sucessão de culturas, comparadas às do campo nativo. *Rev. Bras. Ciênc. Solo* 28, 155–163.
- Bestelmeyer, B.T., 2006. Threshold concepts and their use in rangeland management and restoration: the good, the bad, and the insidious. *Restor. Ecol.* 14, 325–329.
- Boldrini, I.I., 2009. A flora dos Campos do Rio Grande do Sul. In: Pillar, V.D., Müller, S.C., Castilhos, Z.M.S., Jacques, A.V.A. (Eds.), *Campos Sulinos*. MMA, Brasília, pp. 63–77.
- Boldrini, I.I., Eggers, L., 1997. Directionality of succession after grazing exclusion in grassland in the south of Brazil. *Coenoses* 12, 63–66.
- Bond, W.J., Parr, C.L., 2010. Beyond the forest edge: ecology, diversity and conservation of the grassy biomes. *Biol. Conserv.* 143, 2395–2404.
- Brambilla, D.M., et al., 2012. Impact of nitrogen fertilization on the forage characteristics and beef calf performance on native pasture overseeded with ryegrass. *Rev. Brasil. Zootec.* 41, 528–536.
- Briske, D.D., Fuhlendorf, S.D., Smeins, F.E., 2006. A unified framework for assessment and application of ecological thresholds. *Rangeland Ecol. Manage.* 59, 225–236.
- Céspedes-Payret, C., et al., 2012. Land use change in a temperate grassland soil: afforestation effects on chemical properties and their ecological and mineralogical implications. *Sci. Total Environ.* 438, 549–557.
- Cione, A.L., Tonni, E.P., Soibelman, L., 2003. The broken zig-zag: late cenozoic large mammal and tortoise extinction in South America. *Rev. Mus. Argent. Cienc. Nat.* 5, 1–19.
- Cordeiro, J.L.P., Hasenack, H., 2009. Cobertura vegetal atual do Rio Grande do Sul. In: Pillar, V.D., Müller, S.C., Castilhos, Z.M.S., Jacques, A.V.A. (Eds.), *Campos Sulinos*. MMA, Brasília, pp. 285–299.
- Diaz, S., et al., 2007. Plant trait responses to grazing – a global synthesis. *Glob. Change Biol.* 13, 313–341.
- Diekow, J., et al., 2005. Soil C and N stocks as affected by cropping systems and nitrogen fertilisation in a southern Brazil Acrisol managed under no-tillage for 17 years. *Soil Till. Res.* 81, 87–95.
- Favreto, R., et al., 2007. Vegetação espontânea em lavoura sob diferentes manejos estabelecida sobre campo natural. *Iheringia* 62, 5–17.
- Favreto, R., Medeiros, R.B., 2006. Banco de sementes do solo em área agrícola sob diferentes sistemas de manejo estabelecida sobre campo natural. *Rev. Brasil. Sement.* 28, 34–44.
- Fidelis, A., Apuzzato-da-Gloria, B., Pfadenhauer, J., 2009. A importância da biomassa e das estruturas subterrâneas nos Campos Sulinos. In: Pillar, V.D., Müller, S.C., Castilhos, Z.M.S., Jacques, A.V.A. (Eds.), *Campos Sulinos*. MMA, Brasília, pp. 88–100.
- Fidelis, A., et al., 2012. Short-term changes caused by fire and mowing in Brazilian Campos grasslands with different long-term fire histories. *J. Veg. Sci.* 23, 552–562.

- Fidelis, A., et al., 2014. Does disturbance affect bud bank size and belowground structures diversity in Brazilian subtropical grasslands? *Flora* 209, 110–116.
- Fidelis, A., Blanco, C., 2014. Does fire induce flowering in Brazilian subtropical grasslands? *Appl. Veg. Sci.* 17, 690–699.
- Gautreau, P., Velez, E., 2011. Strategies of environmental knowledge production facing land use changes: insights from the Silvicultural Zoning Plan conflict in the Brazilian state of RS. *Cybergeo* 577, <http://cybergeo.revues.org/24881>.
- Gibson, D.J., 2009. *Grasses and grassland ecology*. Oxford, New York.
- Gonçalves, A.R., et al., 2008. Bancos de sementes do sub-bosque de *Pinus spp.* e *Eucalyptus spp.* na Flona de Brasília. *Cerne* 14, 23–32.
- Guo, L., Gifford, R., 2002. Soil carbon stocks and land use change: a meta analysis. *Glob. Change Biol.* 8, 345–360.
- Guo, L.B., et al., 2008. Carbon and nitrogen stocks in a native pasture and an adjacent 16-year-old *Pinus radiata* D. Don. plantation in Australia. *Agric. Ecosyst. Environ.* 124, 205–218.
- Hobbs, R.J., Higgs, E., Harris, J.A., 2009. Novel ecosystems: implications for conservation and restoration. *Trends Ecol. Evol.* 24, 599–605.
- Hobbs, R.J., Higgs, E.S., Hall, C., 2013. *Novel ecosystems: intervening in the new ecological world order*. John Wiley & Sons, Oxford.
- Hölzel, N., et al., 2002. The return of the steppe large-scale restoration of degraded land in southern Russia during the post-Soviet era. *J. Nat. Conserv.* 10, 75–85.
- Hutchings, M.J., Booth, K.D., 1996. Studies on the feasibility of re-creating chalk grassland vegetation on ex-arable land. I. The potential roles of the seed bank and the seed rain. *J. Appl. Ecol.* 33, 1171–1181.
- IBGE - Instituto Brasileiro de Geografia e Estatística, 2012. *Indicadores de desenvolvimento sustentável*. Brasília, IBGE.
- Kiehl, K., Pfadenhauer, J., 2007. Establishment and persistence of target species in newly created calcareous grasslands on former arable fields. *Plant Ecol.* 189, 31–48.
- Lima-Ribeiro, M.S., Diniz-Filho, J.A., 2013. American megafaunal extinctions and human arrival: improved evaluation using a meta-analytical approach. *Quat. Int.* 299, 38–52.
- Mafra, A.F., et al., 2008. Carbono orgânico e atributos químicos do solo em áreas florestais. *Rev. Árvore* 32, 217–224.
- Maia, F.C., et al., 2008. *Lolium multiflorum* seeds in the soil: I. Soil seed bank dynamics in a no till system. *Rev. Brasil. Sement.* 30, 100–110.
- Maraschin, G.E., Corrêa, F.L., 1994. Crescimento e desaparecimento de uma pastagem nativa sob diferentes níveis de oferta de forragem. *Pesq. Agropec. Brasil.* 29, 1617–1623.
- Medeiros, R.B., Pillar, V.D., Reis, J.C.L., 2004. Expansão de *Eragrostis plana* Ness. (Capim Annoni-2) no Rio Grande do Sul e indicativos de controle. *Anales de la 20 Reunión del grupo técnico regional del Cono Sur en mejoramiento y utilización de los recursos forrajeros del área tropical y subtropical, Grupo Campos, Salto, Uruguai*, pp. 208–211.
- Medeiros, R.B., Ferreira, N.R., 2011. Controle de invasão biológica por capim-anomni em margem viária mediante a introdução de gramíneas. *Rev. Brasil. Zootec.* 40, 260–269.
- Meyer, S.T., Koch, G., Weisser, W.W., 2015. Towards a standardized Rapid Ecosystem Function Assessment (REFA). *Trends Ecol. Evol.* 30, 390–397.
- Moojen, E.L., Maraschin, G.E., 2002. Potencial produtivo de uma pastagem nativa do RS submetida a níveis de oferta de forragem. *Ciênc. Rural* 32, 127–132.
- Nabinger, C., et al., 2009. Produção animal com base no campo nativo: aplicações de resultados de pesquisa. In: Pillar, V.D., Müller, S.C., Castilhos, Z.M.S., Jacques, A.V.A. (Eds.), *Campos Sulinos*. MMA, Brasília, pp. 175–198.
- Nabinger, C., Moraes, A., Marschin, G.E., 2000. Campos in Southern Brazil. In: Lemaire, G., Hodgson, J.G., Moraes, A., Nabinger, C., Carvalho, P.C.F. (Eds.), *Grassland ecophysiology and grazing ecology*. CABI, Wallingford, pp. 355–376.
- Öster, M., et al., 2009. Dispersal and establishment limitation reduces the potential for successful restoration of semi-natural grassland communities on former arable fields. *J. Appl. Ecol.* 46, 1266–1274.
- Oliveira, J.M., Pillar, V.D., 2004. Vegetation dynamics on mosaics of Campos and Araucaria forest between 1974 and 1999 in Southern Brazil. *Commun. Ecol.* 5, 197–202.
- Overbeck, G.E., et al., 2005. Fine-scale post-fire dynamics in southern Brazilian subtropical grassland. *J. Veg. Sci.* 16, 655–664.
- Overbeck, G.E., et al., 2007. Brazil's neglected biome: the South Brazilian Campos. *Perspect. Plant Ecol. Evol. Syst.* 9, 101–116.
- Overbeck, G.E., et al., 2013. Restoration ecology in Brazil – time to step out of the forest. *Nat. Conserv.* 11, 92–95.
- Pallarés, O.R., Berretta, E.J., Maraschin, G.E., 2005. The South American campos ecosystem. In: Sutcliffe, J., Reynolds, S.G., Batello, C. (Eds.), *Grasslands of the world*. FAO, Rome, pp. 171–179.
- Perin, E., Ceretta, C., Klamt, E., 2003. Tempo de uso agrícola e propriedades químicas de dois Latossolos do Planalto Médio do RS. *Rev. Bras. Ciênc. Solo* 27, 665–674.
- Pillar, V.D., Boldrini, I., Lange, O., 2002. Spatial patterns of grassland communities under eucalyptus plantation. *Pesq. Agropec. Brasil.* 37, 753–761.
- Pillar, V.D., Tornquist, C.G., Bayer, C., 2012. The southern Brazilian grassland biome: soil carbon stocks, fluxes of greenhouse gases and some options for mitigation. *Braz. J. Biol.* 72, 673–681.
- Pillar, V.D., Vélez, E., 2010. Extinção dos Campos Sulinos em unidades de conservação: um fenômeno natural ou um problema ético? *Nat. Conserv.* 8, 84–86.
- Pinto, C.E., et al., 2008. Produções primária e secundária de uma pastagem natural da Depressão Central do RS submetida a diversas ofertas de fitomassa aérea total. *Rev. Brasil. Zootec.* 37, 1737–1741.
- Podgaiski, L.R., et al., 2013. Spider trait assembly patterns and resilience under fire-induced vegetation change in south Brazilian grasslands. *PLOS ONE* 8, e60207.
- Podgaiski, L.R., et al., 2014. Burning effects on detritivory and litter decay in Campos grasslands. *Aust. Ecol.* 39, 686–695.
- Reed, M.S., et al., 2011. Cross-scale monitoring and assessment of land degradation and sustainable land management: a methodological framework for knowledge management. *Land Degrad. Dev.* 22, 261–271.
- Rheinheimer, D.S., et al., 1998. Modificações em atributos químicos de solo arenoso sob sistema plantio direto. Sociedade Brasileira de Ciência do Solo, Campinas.
- Sala, O.E., et al., 2000. Global biodiversity scenarios for the year 2100. *Science* 287, 1770–1774.
- Sala, O.E., Paruelo, J.M., 1997. Ecosystem services in grasslands. In: Daily, G.C. (Ed.), *Nature's service: Societal dependence on Natural Ecosystems*. Island Press, Washington.
- Schumacher, M.V., Viera, M., Witschoreck, R., 2008. Litter production and nutrients transfer in a second rotation area with *Pinus taeda* L. forest in Cambará do Sul, RS. *Ciênc. Florest.* 18, 471–480.
- Soares, A.B., et al., 2011. Dinâmica da composição botânica numa pastagem natural sob efeito de diferentes ofertas de forragem. *Ciênc. Rural* 41, 1459–1465.
- Souza, A.F., et al., 2013. Afforestation effects on vegetation structure and diversity of grasslands in southern Brazil: the first years. *J. Nat. Conserv.* 21, 56–62.

- Suding, K.N., et al., 2013. Consequences of plant–soil feedbacks in invasion. *J. Ecol.* 101, 298–308.
- Veldman, J.W., et al., 2015. Toward an old-growth concept for grasslands, savannas, and woodlands. *Front. Ecol. Environ.* 13, 154–162.
- Wiesmeier, M., et al., 2009. Depletion of soil organic carbon and nitrogen under *Pinus taeda* plantations in Southern Brazilian grasslands (Campos). *Eur. J. Soil Sci.* 60, 347–359.
- Wilson, J.B., et al., 2012. Plant species richness: the world records. *J. Veg. Sci.* 23, 796–802.
- Winter, S.L., et al., 2012. Restoration of the fire–grazing interaction in *Artemisia filifolia* shrubland. *J. Appl. Ecol.* 49, 242–250.
- Zaloumis, N.P., Bond, W.J., 2011. Grassland restoration after afforestation: no direction home? *Aust. Ecol.* 36, 357–366.
- Zinn, Y.L., Resck, D.V.S., da Silva, J.E., 2002. Soil organic carbon as affected by afforestation with *Eucalyptus* and *Pinus* in the Cerrado region of Brazil. *Forest Ecol. Manage.* 166, 285–294.
- Young, T.P., 2000. Restoration ecology and conservation biology. *Restor. Ecol.* 92, 73–83.