

## Policy Forums

# Imminent threat of the predator fish invasion *Salminus brasiliensis* in a Neotropical ecoregion: eco-vandalism masked as an environmental project



Vinícius Ricardo Ribeiro <sup>a,b,\*</sup>, Pedro Rogério Leandro da Silva <sup>a</sup>, Éder André Gubiani <sup>a,b</sup>, Larissa Faria <sup>c</sup>, Vanessa Salete Daga <sup>c</sup>, Jean Ricardo Simões Vitule <sup>c</sup>

<sup>a</sup> Grupo de Pesquisa em Recursos Pesqueiros e Limnologia (GERPEL), Universidade Estadual do Oeste do Paraná (Unioeste), Toledo, PR, Brazil

<sup>b</sup> Programa de Pós-graduação em Recursos Pesqueiros e Engenharia de Pesca, Universidade Estadual do Oeste do Paraná (Unioeste), Toledo, PR, Brazil

<sup>c</sup> Laboratório de Ecologia e Conservação (LEC), Departamento de Engenharia Ambiental, Setor de Tecnologia, Universidade Federal do Paraná (UFPR), Curitiba, PR, Brazil

## ARTICLE INFO

### Article history:

Received 31 August 2016

Accepted 17 March 2017

Available online 25 May 2017

### Keywords:

Non-native species

Iguaçu basin

Sport fishing

## ABSTRACT

Sport fishing is an activity responsible for moving millions of dollars around the world. However, this activity is also considered to cause habitat degradation and promoting several negative effects on the ichthyofauna. The "Iguaçu Dourado" project, presented here, seeks to intensify the potential of sport fishing in the Iguaçu River Basin. Although not explicit, the project intends to introduce Dourado (*Salminus brasiliensis*), as well as to encourage sport fishing of the species in the Iguaçu River Basin. Twenty specimens were captured in the Salto Santiago and Salto Osório reservoirs from 2003 to 2016, thus indicating that the invasion of *S. brasiliensis* can be successful and extremely harmful to the endemic fauna of the basin. We believe that the so-called project is characterized by an attitude of eco-vandalism, promoting the illegal release of species outside its area of occurrence. Awareness-raising measures as well as educational programs should be addressed so that the population can be informed about the issue of invasion of non-native species.

© 2017 Associação Brasileira de Ciência Ecológica e Conservação. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Sport fishing, introduction of non-native species, biological invasions and their consequences for aquatic ecosystems

Recreational sport fishing is practiced worldwide (Peixer and Petreire Jr., 2009), with resultant important economic and social values in many countries (Pimentel et al., 2001; Britton and Orsi, 2012; Ellender and Weyl, 2014). This fishing modality is responsible for a considerable exchange of income and services related to fishing activities (Sipponen and Gréboval, 2001; Peixer and Petreire Jr., 2009). Despite the positive aspects of sport fishing, related primarily to the economy and to tourism, sport fishing is also considered to be the cause of severe habitat degradation and the decline of fish stocks around the world (e.g. Britton and Orsi, 2012; Ellender and Weyl, 2014), especially by promoting the negative effects of commercial fishing (Cooke and Cowx, 2006).

Presently, sport fishing is one of the main vectors of non-native fish introductions worldwide (Pimentel et al., 2001; Vitule, 2009;

Britton and Orsi, 2012; Daga et al., 2016). In Brazil, on the grounds that few native fish species are suitable for sport fishing, the introduction of non-native species, such as the peacock-bass (*Cichla* spp.) and the dourado *Salminus brasiliensis* (Cuvier, 1816), has been encouraged and even carried out by government agencies, as well as illegally by fishermen, with the aim of developing this fishing modality in the country (Vitule, 2009; Vitule et al., 2014).


The introduction of such species as the dourado and peacock-bass, considered non-native top-predators, can cause significant deleterious effects on native fish populations (Pelicice & Agostinho, 2009; Vitule et al., 2014). Introductions of non-native piscivorous fish can be especially harmful because of their extremely aggressive predatory behavior (Pelicice & Agostinho, 2009; Gubiani et al., 2010; Vitule et al., 2014), and their greater potential for competition when compared with native predators (Pereira et al., 2015).

## The introduction of dourado and prospects for the Iguaçu River basin

Iguaçu River basin is considered to be an unique fish ecoregion (Abell et al., 2008), rich in endemic genera and species

\* Corresponding author.

E-mail address: [viniribeiro\\_23@hotmail.com](mailto:viniribeiro_23@hotmail.com) (V.R. Ribeiro).



**Fig. 1.** Number of individuals of *Salminus brasiliensis* (Cuvier, 1816) recorded per year (bars) and cumulative (line), from 2008 (first record) to 2016, in the Salto Santiago and Salto Osório reservoirs, Iguaçu River basin (A) and sexual maturation stages of the specimens collected in the Iguaçu River basin (imt, immature; mat, maturation; rep, reproduction; spe, spent; res, rest) (B).

(Baumgartner et al., 2012), including many undescribed species (Pavanelli and Bifi, 2009), which is already severely threatened by numerous impoundments and by the introduction of non-native fishes (Gubiani et al., 2010; Daga and Gubiani, 2012; Vitule et al., 2014; Daga et al., 2015, 2016). Currently 29 non-native fish species have been recorded in this region, seven of which are considered top-predators exercising strong predation pressure on native communities (Daga et al., 2016).

The dourado, *Salminus brasiliensis*, is native to southern South America in the Paraná, Paraguay and Uruguay rivers, the Laguna dos Patos drainage, and the Chaparé and Mamoré rivers (Amazon Basin) (Reis et al., 2003; Graça and Pavanelli, 2007). Adults of dourado are aggressive visual predators and primarily piscivorous. The species has recently expanded its range in South America with the help of deliberate introductions (Vitule et al., 2014).

The record of the introduction and subsequent spread of *S. brasiliensis* in some of the main Brazilian river basins, as well as its recent record in the Guaraguaú River basin, is related directly to human activity and, in particular, to sport fishing (Vitule et al., 2014). In the Iguaçu River, the dourado was first recorded in 2008 (Gubiani et al., 2010); records of its occurrence in the river have increased over time (Fig. 1A).

Sampling conducted from 2003 to 2016 in Salto Santiago and Salto Osório reservoirs (for more details see Gubiani et al., 2010; Gerpel/Instituto Neotropical de Pesquisas ambientais/Tractebel, 2015) recorded 20 specimens of dourado (Fig. 1A), with an average standard length of  $40.40 \pm 14.50$  cm (SD) (ranging from 17 to 70 cm) and average weight of  $2106.10 \pm 2359.00$  g (ranging from 88.60 g to 9964.00 g).

The diet of *S. brasiliensis* is predominantly composed of fish, especially species of genus *Astyanax* (Esteves and Pinto Lôbo, 2001). This genus is very abundant and known to be an important food resource for many top-predator fish species in the Iguaçu River (Garavello and Sampaio, 2010; Daga and Gubiani, 2012; Baumgartner et al., 2012; Delariva et al., 2013). In addition, it has a high rate of endemism among its species (Pavanelli and Bifi, 2009). Therefore, because of the high availability of resources, the invasion of *S. brasiliensis* will probably be successful in this region, where it has the potential to be extremely harmful to the endemic fish fauna (Gubiani et al., 2010; Vitule et al., 2014).

The size at first maturity in *S. brasiliensis*, an important measure for inferring the reproductive potential of a fish population (Vazzoler, 1996), was found to range from 37.80 cm for females and 32.40 cm for males (Suzuki et al., 2004). During the period from 2003 to 2016, only two immature males smaller than the recorded size at first maturity were captured, indicating that this species is probably already reproducing in the region (Fig. 1B). However, the migratory behavior of dourado could limit their invasion success

since the area is known to be highly impounded (Daga and Gubiani, 2012; Antonio et al., 2007; Daga et al., 2015). Nevertheless, the invasion success of *S. brasiliensis* in other Neotropical rivers with similar limitations on migration has already been reported (Ruschi, 1965; Vitule et al., 2014), which leads us to conclude that its successful establishment in the Iguaçu River is only a matter of time.

#### Project 'Iguaçu Dourado' and the risks associated with the introduction of the dourado in the Iguaçu River basin

The project 'Iguaçu Dourado' (Fig. 2A) aims at the implementation of a program for the sustainable exploitation of sport fishing tourism in the Iguaçu River basin. In its publicity material the project claims to be in favor of the conservation of the region and the sustainable development of the local population (Fig. 2B). Although not explicit, the intention of the group is to introduce and/or maintain the dourado by catch and release programs, as well as by clandestine restocking by individual or group activity, is clear (e.g. Fig. 2C and D), even though the organizers are aware that this practice is prohibited by the federal laws nos. No. 5197/67 and No. 9605/98. Therefore, this project can be characterized as embodying an attitude of eco-vandalism, and can be understood as the deliberate and illegal release of species outside of their natural areas of occurrence (Cambray, 2003; Abilhoa et al., 2011). In addition, one of the most negative points of Project 'Iguaçu Dourado' is that it is masked as a correct environmental initiative (see attached links on capture and restocking of non-native species in the Iguaçu River basin). According to Azevedo-Santos et al. (2015), environmental education and information are the best ways to avoid the introduction of fish species, but in Project 'Iguaçu Dourado' they are used to abet a completely wrong initiative. The project's offer of "educational" workshops, especially those for children from six to 12 years old, is, in reality, an attempt to induce children and young people to think that what they are doing is correct, which completely undermines the environmental education of future citizens.

Therefore, we wish to emphasize strongly that this method of promoting sustainable sport fishing tourism in the Iguaçu River is inappropriate, especially since it implicitly suggests the use of a non-native species. In order to achieve the real purposes of this project, the development of sustainable sport fishing in the region, the incentive for sport fishing must be directed toward native fish species, and combined with an educational awareness program and with sustainable recreational sport fishing practices, under the supervision of the appropriate public agencies (e.g. Azevedo-Santos et al., 2015).

Legal measures to prevent the introduction of non-native species in Brazil have failed so far, precisely due to the lack of governmental supervision and the lack of public information about

**A Iguacu Dourado**

Programa para Exploração Sustentável do Turismo de Pesca do Rio Iguacu

**C 3. DESCRIÇÃO**

A realização de estudos voltados ao diagnóstico quantitativo e qualitativo para manejo da ictiofauna em áreas previamente selecionadas para a conservação ambiental.

Fomento da atividade de turismo de pesca sustentável nos locais determinados como "áreas-piloto".

Treinamento, capacitação e sensibilização das comunidades e profissionais envolvidos na atividade.

**B INTRODUÇÃO**

Conforme tendências internacionais, o progresso ligado à conservação do meio ambiente não só é um bom investimento como também uma solução viável aos problemas sociais de regiões onde os atrativos naturais estejam conservados.

Símbolo de toda uma região, o rio Iguacu merece uma atenção especial pelo seu potencial. Além da energia já produzida pela suas hidrelétricas, outras riquezas se escondem em suas águas. O vale do Médio Iguacu tem vocação para tirar do turismo da pesca suporte para a almejada sustentabilidade e este Projeto pretende dar subsídios para a exploração deste potencial.

**D 2. JUSTIFICATIVA**

Apesar das características do Médio Rio Iguacu se enquadrem no perfil de área com vocação para explorar atrativos turísticos, suas qualidades não estão sendo aproveitadas.

Com área de influência em vários Municípios, o ecoturismo pode gerar receita considerável, com atividades capazes de substituir o extrativismo danoso ao ambiente.

A riqueza em espécies esportivas e uma área com variados ambientes comprovam ter a Pesca Amadora potencial para ser um grande chamariz ao ecoturismo.

Para citarmos o potencial e as dimensões desta atividade, o impacto anual gerado pela pesca esportiva na vizinha Argentina, em rios com características semelhantes, é um cifra que gira na casa dos milhões de dólares. Comparando as realidades vividas em nossas águas, a exploração de nosso potencial está aquém do possível e viável.

Os Estados do Paraná e Santa Catarina contam com grau de desenvolvimento na Pesca Esportiva destacado em relação aos demais Estados do Brasil. Preenchemos todos os pré requisitos para nos tornarmos referência nesta atividade que, infelizmente só está sendo incentivada em outras regiões.

Para melhor representatividade dos resultados e otimização de recursos investidos, sugerimos a adoção de Projeto Piloto numa área restrita onde as ações teriam início, para posterior maior abrangência.

As etapas seguintes teriam adequações às características individuais da região visada.

Entre as opções, a região situada entre o Município de Porto Amazonas, a montante e, a jusante, o Município de Saudades do Iguaçu contém áreas com grau de conservação satisfatório, acesso viável, delimitação racional, fiscalização carente, com variedade de peixes esportivos e infraestrutura possível de ser adaptada para atender o turista. Portanto são locais com vocação para implantarmos nosso projeto, abrangendo seu todo ou em regiões segmentadas

**Fig. 2.** Publicity material of project for the sustainable exploitation of sport fishing tourism in the Iguaçu River basin, entitled 'Iguacu Dourado': cover material (A), project introduction (B), description of the proposal for the development of sport fishing in the basin (C) and justification for the project (D). Text translation into English is in Supplementary Material.

the existence of specific legislation (Vitule, 2009; Azevedo-Santos et al., 2015). What is needed is a change in the behavior of the population through education and information, the most effective way to mitigate the deliberate introductions of non-native species (Azevedo-Santos et al., 2015). Additionally, the media in general, and more specifically TV shows and magazines about sport fishing, should act as channels of information concerning which species are not native to a region, and be aimed at informing and educating fishermen, so that they can become active partners in combating the invasion of non-native species (Vitule, 2009).

#### Conflicts of interest

The authors declare no conflicts of interest.

#### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.pecon.2017.03.004.

#### References

- Abell, R., Thieme, M.L., Revenga, C., 2008. Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience* 58, 403–414.
- Abilio, V., Braga, R.R., Bornatowski, H., et al., 2011. Fishes of the Atlantic Rain Forest streams: ecological patterns and conservation, changing diversity in changing environment. In: Grillo, O., Venora, G. (Eds.), *Changing Diversity in Changing Environment*. InTech., pp. 259–282, Available from: <http://www.intechopen.com/books/changing-diversity-in-changing-environment/fishes-of-the-atlantic-rainforest-streams--ecological-patterns-and-conservation>.
- Antonio, R.R., Agostinho, A.A., Pelicice, F.M., 2007. Blockage of migration routes by dam construction: can migratory fish find alternative routes? *Neotrop. Ichthyol.* 5 (2), 177–184.
- Azevedo-Santos, V.M., Pelicice, F.M., Lima-Junior, D.P., 2015. How to avoid fish introductions in Brazil: education and information as alternatives. *Natur. Conserv.* 13, 123–132.
- Baumgartner, G., Pavanelli, C.S., Baumgartner, D., et al., 2012. Peixes do Baixo Rio Iguaçu. *Eduem, Maringá*.
- Britton, J.R., Orsi, M.L., 2012. Non-native fish in aquaculture and sport fishing in Brazil: economic benefits versus risks to fish diversity in the upper River Paraná Basin. *Rev. Fish Biol. Fish.* 22, 555–565.
- Cambray, J.A., 2003. Impact on indigenous species biodiversity caused by the globalisation of alien recreational freshwater fisheries. *Hydrobiologia* 500, 217–230.

- Cooke, S., Cowx, I.G., 2006. Contrasting recreational and commercial fishing: searching for common issues to promote unified conservation of fisheries resources and aquatic environments. *Biol. Conserv.* 128, 93–108.
- Daga, V.S., Gubiani, E.A., 2012. Variations in the endemic fish assemblage of a global freshwater ecoregion: associations with introduced species in cascading reservoirs. *Acta Oecol.* 41, 95–105.
- Daga, V.S., Skora, F., Padial, A.A., 2015. Homogenization dynamics of the fish assemblages in Neotropical reservoirs: comparing the roles of introduced species and their vectors. *Hydrobiologia* 746, 327–347.
- Daga, V.S., Debona, T., Abilhoa, V., 2016. Non-native fish invasions of a Neotropical ecoregion with high endemism: a review of the Iguaçu River. *Aquat. Invas.* 11, 209–223.
- Delariva, R.L., Hahn, N.S., Kashiwaqui, E.A.L., 2013. Diet and trophic structure of the fish fauna in a subtropical ecosystem: impoundment effects. *Neotrop. Ichthyol.* 11 (4), 891–904.
- Ellender, B., Weyl, O.L.F., 2014. A review of current knowledge, risk and ecological impacts associated with non-native freshwater fish introductions in South Africa. *Aquat. Invas.* 9, 117–132.
- Esteves, K.E., Pinto Lôbo, A.V., 2001. Feeding pattern of *Salminus maxillosus* (Pisces Characidae) at Cachoeira das Emas, Mogi-Guaçu River (São Paulo State Southeast Brazil). *Braz. J. Biol.* 61 (2), 267–276.
- Garavello, J.C., Sampaio, F.A.A., 2010. Five new species of genus *Astyanax* Baird & Girard, 1854 from Rio Iguaçu, Paraná, Brazil (Ostariophysi Characiformes, Characidae). *Braz. J. Biol.* 70, 847–865.
- Gerpel/Istituto Neotropical de Pesquisas ambientais/Tractebel., 2015. Estudos ictiológicos e monitoramento da qualidade de água dos reservatórios de Salto Santiago e Salto Osório – Rio Iguaçu. 180p.
- Graça, W.J., Pavanello, C.S., 2007. Peixes da planície de inundação do alto rio Paraná e áreas adjacentes. Eduem, Maringá.
- Gubiani, E.A., Frana, V.A., Maciel, A.L., 2010. Occurrence of the non-native fish *Salminus brasiliensis* (Cuvier, 1816), in a global biodiversity ecoregion, Iguaçu River, Paraná River basin, Brazil. *Aquat. Invas.* 5, 223–227.
- Pavanelli, C.S., Bifi, A.G., 2009. A new *Tatia* (Ostariophysi: Siluriformes: Auchenipteridae) from the rio Iguaçu basin, Paraná State Brazil. *Neotrop. Ichthyol.* 7 (2), 199–204.
- Peixer, J., Petreiro Jr., M., 2009. Sport fishing in Cachoeira de Emas in Mogi-Guaçu River, State of São Paulo, Brazil. *Braz. J. Biol.* 69, 1081–1090.
- Pelicice, F.M., Agostinho, A.A., 2009. Fish fauna destruction after the introduction of non-native predator (*Cichla kelberi*) in a neotropical reservoir. *Biol. Invas.* 11, 1789–1801.
- Pereira, L.S., Agostinho, A.A., Gomes, L.C., 2015. Eating the competitor: a mechanism of invasion. *Hydrobiologia* 746, 223–231.
- Pimentel, D., Macknair, S., Janecka, J., 2001. Economic and environmental threats of alien plant, animal, and microbe invasions. *Agricultr. Ecosyst. Environ.* 84, 1–20.
- Reis, R.E., Kullander, S.O., Ferraris-Jr, C.F., 2003. Check list of the freshwater fishes of South and Central America. Editora PUCRS, Porto Alegre.
- Ruschi, A., 1965. Lista dos tubarões, raias e peixes de água doce e salgada do estado do Espírito Santo e uma observação sobre a introdução do dourado no rio Doce. *Boletim Museu Biol. Mello Leitão* 25A, 1–23.
- Sipponen, M., Gréboval, D., 2001. Social, economic and cultural perspectives of European inland fisheries: review of the EIFAC symposium on fisheries and society. *Fish. Manage. Ecol.* 8, 283–293.
- Suzuki, H.I., Vazzoler, A.E.A.M., Marques, E.E., 2004. Reproductive ecology of the fish assemblages. In: Thomaz, S.M., Agostinho, A.A., Hahn, N.S. (Eds.), The Upper Paraná River and its Floodplain: Physical Aspects, Ecology and Conservation. Backhuys Publishers, Leiden, pp. 271–291.
- Vazzoler, A.E.A.M., 1996. Biologia da reprodução de peixes teleósteos: teoria e prática. Eduem, Maringá.
- Vitule, J.R.S., 2009. Introdução de peixes em ecossistemas continentais brasileiros: revisão, comentários e sugestões de ações contra o inimigo quase invisível. *Neotrop. Bio. Conserv.* 4 (2), 111–122.
- Vitule, J.R.S., Bornatowski, H., Freire, C.A., 2014. Extralimital introductions of *Salminus brasiliensis* (Cuvier, 1816) (Teleostei, Characidae) for sport fishing purposes: a growing challenge for the conservation of biodiversity in neotropical aquatic ecosystems. *BiolInvas. Rec.* 3 (4), 291–296.